

# Search for the quark shell structure using the non-topological soliton model

N.Sawado<sup>\*</sup>, N.Shiiki and S.Oryu

*Department of Physics, Tokyo University of Science,  
2641, Yamazaki, Noda, Chiba 278-8510, Japan*

## Abstract

We investigate higher baryon-number states in the Chiral Quark Soliton Model using the rational map ansatz for the background chiral fields. The soliton solutions are obtained self-consistently. We show that the baryon number density has point-like symmetries and the corresponding valence quark spectrum of the lowest energy exhibits approximate four-fold degenerate. Our results indicate the possibility of existence of the shell-like structure in the multi-baryonic system.

PACS numbers: 12.39.Fe, 12.39.Ki, 21.60.-n, 24.85.+p

---

<sup>\*</sup> sawado@ph.noda.tus.ac.jp

The Chiral Quark Soliton Model (CQSM) was developed in 1980's as an effective theory of QCD interpolating between the Constituent Quark Model and Skyrme Model [1, 2]. In the large  $N_c$  limit, these models are identical [3].

The CQSM is derived from the instanton liquid model of the QCD vacuum and incorporates the non-perturbative feature of the low-energy QCD, spontaneous chiral symmetry breaking. The vacuum functional is defined by;

$$\mathcal{Z} = \int \mathcal{D}\pi \mathcal{D}\psi \mathcal{D}\psi^\dagger \exp \left[ i \int d^4x \bar{\psi} (i\not{\partial} - mU^{\gamma_5}) \psi \right] \quad (1)$$

where the SU(2) matrix

$$U^{\gamma_5} = \frac{1 + \gamma_5}{2} U + \frac{1 - \gamma_5}{2} U^\dagger \quad \text{with} \quad U = \exp(i\vec{\tau} \cdot \vec{\pi}/f_\pi)$$

describes chiral fields,  $\psi$  is quark fields and  $m$  is the dynamical quark mass.  $f_\pi$  is the pion decay constant and experimentally  $f_\pi \sim 93\text{MeV}$ . Since our concern is the tree-level pions and one-loop quarks according to the Hartree mean field approach, the kinetic term of the pion fields which gives a contribution to higher loops can be neglected. Due to the interaction between the valence quarks and the Dirac sea, soliton solutions appear as bound states of quarks in the background of self-consistent mean chiral field.  $N_c$  valence quarks fill the each bound state to form a baryon. The baryon number is thus identified with the number of bound states filled by the valence quarks [4]. For  $B = 1$  and 2, the spherically symmetric soliton [5, 6] and the axially symmetric soliton [7] were found respectively. Upon quantization, the intermediate states of nucleon and deuteron between the Constituent Quark Model and Skyrme Model were obtained.

The vacuum functional in Eq.(1) can be integrated over the quark fields to obtain the sea quark energy [8]

$$E_{sea}[U] = \frac{1}{4\sqrt{\pi}} N_c \int_{1/\Lambda^2}^{\infty} \frac{d\tau}{\tau^{3/2}} \left( \sum_{\nu} e^{-\tau\epsilon_{\nu}^2} - \sum_{\nu} e^{-\tau\epsilon_{\nu}^{(0)2}} \right) . \quad (2)$$

where  $\Lambda$  is cut-off parameter in the proper-time regularization [9] and  $\epsilon_{\nu}, \epsilon_{\nu}^{(0)}$  are the eigenvalues of Dirac equation for a single quark with the vacuum sector defined by

$$\left( \frac{\vec{\alpha} \cdot \vec{\nabla}}{i} + \beta m U^{\gamma_5} \right) |\nu\rangle = \epsilon_{\nu} |\nu\rangle , \quad \left( \frac{\vec{\alpha} \cdot \vec{\nabla}}{i} + \beta m \right) |\nu\rangle^{(0)} = \epsilon_{\nu}^{(0)} |\nu\rangle^{(0)} . \quad (3)$$

In the Hartree picture, the baryon states are the quarks occupying all negative Dirac sea and valence levels. Hence, if we define the total soliton energy  $E_{total}$ , the valence quark energy should be added;

$$E_{total}[U] = N_c \sum_i E_{val}^{(i)}[U] + E_{sea}[U] . \quad (4)$$

where  $E_{val}^{(i)}$  is the valence quark contribution to the  $i$  th baryon.

For higher winding number solitons, it is expected the solution to have point-like symmetries from the study of the Skyrme model [10]. Therefore, we shall impose the same symmetries on the chiral fields using the rational map ansatz. According to the ansatz, the chiral field can be expressed as [11]

$$U(r, z) = \exp (if(r)\vec{n}_R \cdot \vec{\tau}) \quad (5)$$

where

$$\vec{n}_R = \frac{1}{1 + |R(z)|^2} (2\text{Re}[R(z)], 2\text{Im}[R(z)], 1 - |R(z)|^2)$$

and  $R(z)$  is a rational map.

Rational maps are maps from  $CP(1)$  to  $CP(1)$  (equivalently, from  $S^2$  to  $S^2$ ) classified by winding number. It was shown in [11] that  $B = N$  skyrmions can be well-approximated by rational maps with winding number  $N$ . The rational map with winding number  $N$  possesses  $(2N + 1)$  complex parameters whose values can be determined by imposing the symmetry of the skyrmion. Thus, the rational maps for  $B = 2 \sim 7$  and  $B = 17$  take the form;

$$\begin{aligned} R(z) &= z^2 \ (B = 2) , \ R(z) = \frac{\sqrt{3}iz^2 - 1}{z(z^2 - \sqrt{3}i)} \ (B = 3) , \\ R(z) &= \frac{z^4 + 2\sqrt{3}iz^2 + 1}{z^4 - 2\sqrt{3}iz^2 + 1} \ (B = 4) , \\ R(z) &= \frac{z(z^4 + 3.94z^2 + 3.07)}{3.07z^4 - 3.94z^2 + 1} \ (B = 5) , \ R(z) = \frac{z(z^4 - 5)}{-5z^4 + 1} \ (B = 5^*) , \\ R(z) &= \frac{z^4 + 0.16i}{z^2(0.16z^4i - 1)} \ (B = 6) , \\ R(z) &= \frac{7/\sqrt{5}z^6 - 7z^4 - 7/\sqrt{5}z^2 - 1}{z(z^6 + 7/\sqrt{5}z^4 + 7z^2 - 7/\sqrt{5})} \ (B = 7) , \\ R(z) &= \frac{17z^{15} - 187z^{10} + 119z^5 - 1}{z^2(z^{15} + 119z^{10} + 187z^5 + 17)} \ (B = 17) , \end{aligned} \quad (6)$$

where the complex coordinate  $z$  on  $CP(1)$  is identified with the polar coordinates on  $S^2$  by  $z = \tan(\theta/2)e^{i\varphi}$  via stereographic projection. Substituting (6) into (5), one obtains the complete form of the chiral fields with appropriate symmetry and winding number. Since the chiral fields in (5) are parameterized by the polar coordinates, one can apply the numerical technique developed for  $B = 1$  to find higher soliton solutions [6].

Demanding that the total energy in (4) be stationary with respect to variation of the profile function  $f(r)$ ,

$$\frac{\delta}{\delta f(r)} E_{total} = 0 ,$$

yields the field equation

$$S(r) \sin f(r) = P(r) \cos f(r) \quad (7)$$

where

$$S(r) = N_c \sum_{\nu} (n_{\nu} \theta(\epsilon_{\nu}) + \text{sign}(\epsilon_{\nu}) \mathcal{N}(\epsilon_{\nu})) \langle \nu | \gamma^0 \delta(|x| - r) | \nu \rangle \quad (8)$$

$$P(r) = N_c \sum_{\nu} (n_{\nu} \theta(\epsilon_{\nu}) + \text{sign}(\epsilon_{\nu}) \mathcal{N}(\epsilon_{\nu})) \langle \nu | i \gamma^0 \gamma^5 \vec{n}_R \cdot \vec{\tau} \delta(|x| - r) | \nu \rangle . \quad (9)$$

The procedure to obtain the self-consistent solution of Eq.(7) is that 1) solve the eigenequation in (3) under an assumed initial profile function  $f_0(r)$ , 2) use the resultant eigenfunctions and eigenvalues to calculate  $S(r)$  and  $P(r)$ , 3) solve Eq.(7) to obtain a new profile function, 4) repeat 1) – 3) until the self-consistency is attained.

Fig. 1 ~ Fig. 5 shows the spectrums of the quark orbits for  $B = 3 \sim 7$  as a function of the soliton-size parameter  $X$ . In  $B = 3$ , the  $P$  orbit diving into the negative energy region is triply degenerate. As discussed in [4], baryon number of the soliton equals to the number of diving levels occupied by  $N_c$  valence quarks. Thus putting  $N_c = 3$  valence quarks on each of the degenerate levels, one obtains the  $B = 3$  soliton solution. Similarly, for higher  $B$  than 3, the same number of orbits as  $B$  dive into negative energy region.

Fig. 6 and Fig. 7 show the corresponding baryon density and profile functions. This result is consistent with the result for skyrmions obtained by Braaten *et.al* [10] and Houghton *et.al*. [11].

In Fig. 8, the valence quark spectra are shown for  $B = 1 \sim 7$  and  $B = 17$ . The approximate four-fold degeneracy can be observed for the lowest level. However, in the

CQSM, a quark has only two quantum numbers: colour and grand spin defined by  $\vec{K} = \vec{L} + \frac{1}{2}\vec{\sigma} + \frac{1}{2}\vec{\tau}$ , which means the lowest level ( $K = 0$ ) should show no degeneracy. At present, all we can say is that if one employs point-like symmetries for the background chiral fields, one obtains the larger degeneracy of a single quark states than the spherical symmetric case. Consequently, the number of the states can be reduced and the large shell gaps emerge. These results indicate the possibility of the existence of the lowest energy shell filled up with four quarks and also, these may be the reason why  ${}^4He$  is stable compared with other nuclei.

### Acknowledgements

We are grateful to N.S.Manton for suggesting us this subject and the possibility of the shell structure for quarks.

- 
- [1] D. I. Diakonov and V. Yu. Petrov, Nucl. Phys. **B306**, 457 (1986).
  - [2] R. Alkofer, H. Reinhardt and H. Weigel, Phys. Rept. **265**, 139 (1996); Chr. V. Christov, A. Blotz, H.-C.Kim, P. Pobylitsa, T. Watabe, Th. Meissner, E. Ruiz Arriola, K. Goeke, Prog. Part. Nucl. Phys. **37**, 91 (1996).
  - [3] A. Manohar, Nucl. Phys. **B248**, 19 (1984).
  - [4] S. Kahana and G. Ripka, Nucl. Phys. **A429**, 462 (1984).
  - [5] H. Reinhardt and R. Wünsch, Phys. Lett. **B215**, 577 (1988).
  - [6] M. Wakamatsu and H. Yoshiki, Nucl. Phys. **A524**, 561 (1991).
  - [7] N. Sawado and S. Oryu, Phys. Rev. **C58**, R3046 (1998); N. Sawado, Phys. Rev. **C61**, 65206 (2000).
  - [8] Th. Meissner, E. Ruiz Arriola, F. Grümmer, H. Mavromatis and K. Goeke, Phys. Lett. **B214**, 312 (1988).
  - [9] J. Schwinger, Phys. Rev. **82**, 664 (1951).
  - [10] E. Braaten, S. Townsend and L. Carson, Phys. Lett. **B235**, 147 (1990).
  - [11] C. J. Houghton, N. S. Manton and P. M. Sutcliffe, Nucl. Phys. **B510**, 507 (1998).

TABLE I: Spectrums of the valence, vacuum sea and their sum (in MeV).  $\bar{M}$  is defined as  $\bar{M} \equiv M/BM_{B=1}$  which means the energy per baryon. The results for  $\bar{M}$  in Skyrme model calculations [11] are also quoted.

| $B$ | Valence |     |     |     |     |     |     | Sea  | Total | $\bar{M}$ (Skyrme) |
|-----|---------|-----|-----|-----|-----|-----|-----|------|-------|--------------------|
| 1   | 173     |     |     |     |     |     |     | 674  | 1192  | 1.00 (1.00)        |
| 3   | 210     | 210 | 210 |     |     |     |     | 1554 | 3444  | 0.96 (0.96)        |
| 4   | 153     | 156 | 156 | 156 |     |     |     | 2544 | 4407  | 0.92 (0.92)        |
| 5   | 123     | 131 | 131 | 139 | 210 |     |     | 3265 | 5467  | 0.92 (0.93)        |
| 5*  | 157     | 157 | 157 | 232 | 232 |     |     | 2874 | 5680  | 0.95 (1.00)        |
| 7   | 115     | 120 | 120 | 120 | 166 | 166 | 166 | 4554 | 7478  | 0.90 (0.90)        |




FIG. 1: Spectrum of the quark orbits for  $B = 3$  are illustrated as a function of the “soliton size”  $X$ . Trial profile function is defined by  $f(r) = -\pi + \pi r/X$  for  $X \leq r$  and  $f(r) = 0$  for  $X > r$ .




FIG. 2: Spectrum of the quark orbits for  $B = 4$ .






FIG. 3: Spectrum of the quark orbits for  $B = 5$   $D_{2d}$  symmetry.




FIG. 4: Spectrum of the quark orbits for  $B = 5$  octahedron.




FIG. 5: Spectrum of the quark orbits for  $B = 7$  dodecahedron.




FIG. 6: Surface of the baryon-number density: (a)  $B = 3$  tetrahedron, (b)  $B = 4$  cube, (c)  $B = 5$  with  $D_{2d}$  symmetry, (d)  $B = 5$  octahedron, (e)  $B = 7$  dodecahedron.




FIG. 7: Profile functions  $f(r)$  of the rational map ansatz.




FIG. 8: Valence quark spectrums of  $B = 1 \sim 7$  and  $B = 17$ .